

## ABSTRACT

### The hemostatic pathway in ischemic stroke

Clinical studies of genetic variation and plasma protein measurements

Ellen Hanson

Department of Clinical Neuroscience and Rehabilitation, Institute of Neuroscience and  
Physiology, University of Gothenburg, Göteborg, Sweden, 2012

Although stroke is a common cause of death and disability in adults, there are few studies on stroke compared to other common diseases. A stroke could be either ischemic or hemorrhagic, but even within these two groups, disease etiology shows heterogeneity. In ischemic stroke, the different etiologic subtypes represent different underlying pathophysiologic mechanisms. However, the formation of a thrombus is a key mechanistic event in the majority of ischemic stroke events. Therefore, the aim of the present thesis was to test the hypothesis that hemostatic gene polymorphisms and/or plasma levels of hemostatic proteins are associated with ischemic stroke. A second aim was to investigate whether the associations differ between the etiologic subtypes of ischemic stroke.

The studies were based on the Sahlgrenska Academy Study on Ischemic Stroke (SAHL SIS), which includes 844 patients with ischemic stroke and 668 controls, all younger than 70 years of age. Patients were classified into the major etiologic subtypes of ischemic stroke, *i.e.* large-vessel disease, small-vessel disease, cardioembolic stroke and cryptogenic stroke. Genotyping was carried out using both low- and high-throughput methods. Plasma levels of hemostatic proteins were determined by immunological methods.

The initial studies in this thesis focused the von Willebrand factor (VWF) as well as the VWF-cleaving protease ADAMTS13. We found that ADAMTS13 gene variation was associated with overall ischemic stroke and with the etiologic subtype of cryptogenic stroke. Regarding VWF, the plasma levels were increased in overall ischemic stroke, as well as in all four major etiologic subtypes, as compared to the controls. There were also significant differences in VWF levels between the subtypes, highlighting the importance of considering etiologic subtypes in ischemic stroke studies. ABO blood group strongly influences VWF plasma levels, but we found no association between ABO and ischemic stroke. We then went on by analyzing plasma levels and gene variants of the newly discovered factor VII-activating protease (FSAP). FSAP gene variation influenced the plasma levels, but was not associated with ischemic stroke. Plasma FSAP on the other hand, was independently associated with overall ischemic stroke and with all major etiologic subtypes, indicating that FSAP is involved in ischemic stroke independent of the underlying etiology. We also observed an association between coagulation factor XI (FXI) gene variants and overall ischemic stroke up to 70 years of age, suggesting that FXI might be involved in ischemic stroke with a relatively young age of onset.

In conclusion, these results support a role for prothrombotic mechanisms in the pathophysiology of ischemic stroke. These mechanisms appear to be of importance for all four major etiologic subtypes of ischemic stroke, while we also show that there are subtype-specific differences.

**Keywords:** ischemic stroke, etiologic subtypes of ischemic stroke, genetics, SNP, hemostasis, prothrombotic, ADAMTS13, VWF, ABO blood group, FSAP, FXI

ISBN 978-91-628-8518-2

# The hemostatic pathway in ischemic stroke

Clinical studies of genetic variation and plasma protein measurements

Akademisk avhandling

som för avläggande av medicine doktorsexamen vid Sahlgrenska akademien  
vid Göteborgs universitet kommer att offentligen försvaras i hörsal  
Arvid Carlsson, Academicum, Medicinaregatan 3  
fredagen den 14 september 2012 kl. 9.00

av Ellen Hanson

Fakultetsopponent: Professor Jan-Håkan Jansson  
Folkhälsa och klinisk medicin, Umeå universitet

- I. **Ellen Hanson**, Katarina Jood, Staffan Nilsson, Christian Blomstrand, Christina Jern. Association between genetic variation at the ADAMTS13 locus and ischemic stroke. *Journal of Thrombosis and Haemostasis* 2009 Dec;7(12):2147-49.
- II. **Ellen Hanson**, Katarina Jood, Sara Karlsson, Staffan Nilsson, Christian Blomstrand, Christina Jern. Plasma von Willebrand factor in the etiologic subtypes of ischemic stroke. *Journal of Thrombosis and Haemostasis* 2011 Feb;9(2):275-81.
- III. **Ellen Hanson**,\* Sara Karlsson,\* Katarina Jood, Staffan Nilsson, Christian Blomstrand, Christina Jern. No evidence for an association between ABO blood group and overall ischemic stroke or any of the etiologic subtypes. *Thrombosis Research* 2012 Apr 4. *In press*
- IV. **Ellen Hanson**,\* Sandip M Kanse,\* Amit Joshi, Staffan Nilsson, Katarina Jood, Christian Blomstrand, Christina Jern. Plasma factor VII-activating protease antigen levels and activity are increased in ischemic stroke. *Journal of Thrombosis and Haemostasis* 2012 May;10(5):848-56.
- V. **Ellen Hanson**, Staffan Nilsson, Katarina Jood, Bo Norrving, Gunnar Engström, Christian Blomstrand, Arne Lindgren, Olle Melander, Christina Jern. Genetic variants of coagulation factor XI show association with ischemic stroke up to 70 years of age. *In manuscript*

\*Both authors contributed equally to this work.

Göteborg 2012



UNIVERSITY OF GOTHENBURG